

# **Committee Advice on the safety of cannabidiol (CBD) isolate as a novel food for use in food supplements - RP354**

## **Reference number RP354**

Food Standards Agency (FSA) and Food Standards Scotland (FSS)

Regulated Product Dossier Assessment

Assessment finalised: 18th of February 2025

## **Summary**

An application was submitted to the Food Standards Agency (FSA) and Food Standards Scotland (FSS) in February 2021 from Bridge Farm Group. (“the applicant”) for the authorisation of cannabidiol (CBD) isolate as a novel food.

The novel food is a CBD isolate which is intended to be used as a food ingredient in food supplements for adults (excluding pregnant and lactating women and other specifically identified vulnerable groups, including those taking medication and the immunosuppressed).

The novel food was assessed based on the data provided. This review indicated it was appropriate for the provisional ADI for 98% or greater CBD to form part of the evidence for this assessment. For CBD a provisional acceptable daily intake (ADI) of 10 mg/day for a healthy 70kg adult has been published by the FSA and was considered in assessing this novel food. The provisional ADI (section 2.7) was recommended, subject to the existing advice to consumers that pregnant and breastfeeding women and people taking any prescription medication should avoid the consumption of CBD. Consumers on regular medications should seek advice from a medical professional before using any type of CBD food product. In addition, children and prospective parents trying for a baby are advised against

consumption of CBD, as are those who are immunosuppressed, due to remaining data gaps and residual uncertainties concerning the safety of CBD for these groups of consumers. These contraindications would also apply to this novel food.

To support the FSA and FSS in their evaluation of the application, the Advisory Committee on Novel Foods and Processes (ACNFP) were asked to review the safety dossier and supplementary information provided by the applicant. The Committee did not consider any potential health benefits or claims arising from consuming the food, as the focus of the novel food assessment is to ensure the food is safe and does not put consumers at a nutritional disadvantage.

The Committee concluded that the applicant had provided sufficient information to assure the novel food, an isolated CBD as detailed in application RP 354, was safe under the proposed conditions of use. The anticipated intake levels and the proposed use in food supplements was not considered to be nutritionally disadvantageous.

## **1. Introduction**

1. The ACNFP assessed the food safety risks of CBD isolated from hemp (*Cannabis sativa*) and its production under the proposed uses in line with Article 7 of assimilated Commission Implementing Regulation (EU) 2017/2469. The regulatory framework and the technical guidance put in place by the European Food Safety Authority (EFSA) for full novel food applications is applicable and formed the basis and structure for the assessment (EFSA NDA Panel, 2021).

2. An application was submitted to the Food Standards Agency (FSA) and Food Standards Scotland (FSS) in February 2021 from Bridge Farm Group. ("the applicant") for the authorisation of isolated cannabidiol (CBD) as described in RP 354, as a novel food. The novel food is a >98% pure, hemp-derived CBD isolate, which is intended to be used as an ingredient in food supplements for adults (excluding pregnant and lactating women and other specifically identified vulnerable groups including those taking medication and the immunosuppressed).

3. Advice was sought from the joint Subgroup of the ACNFP and the Committee on Toxicity (COT) on CBD and hemp derived products on the quality of the toxicological evidence submitted to support the application. The ACNFP and COT have issued a joint statement on the safe upper intake of ingredients containing 98% or more CBD. This was considered in reviewing the toxicological evidence for this application.

4. Following the review by the ACNFP at the 169<sup>th</sup> final recommendations from the Committee were presented, allowing the Committee Advice to be concluded.
5. This document outlines the conclusions of the ACNFP on the safety of a cannabidiol (CBD) isolate as detailed in application RP 354, as a novel food.

## **2. Assessment**

### **2.1 Identity of novel food**

6. The novel food is a Cannabidiol (CBD) isolate in the form of an amorphous powder of purity equal to or greater than 98.0%. Information to support this characterisation was provided for five batches of the novel food.
7. CBD is characterised by the chemical formula: C<sub>21</sub>H<sub>30</sub>O<sub>2</sub>; molecular mass: 314.46 g/mol; CAS number: 13956-29-1; IUPAC name: 2-[(1R,6R)-6-Isopropenyl-3-methylcyclohex-2-en-1-yl]-5-pentylbenzene-1,3-diol.

#### **Figure 1: The molecular structure of CBD.**

8. Confirmation of its identity and purity was provided by Quantitative Nuclear Magnetic Resonance Spectroscopy q-(NMR) and Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS).

### **2.2 Production Process**

9. The CBD isolate is manufactured using a multi-step process under controlled conditions.
10. Certificates of analysis for raw starting materials used in the extraction process were provided to demonstrate the effectiveness of the controls at this point in the process. The details of the commercially sensitive extraction process were shared and reviewed by the ACNFP.
11. Plants are grown under conditions of good horticultural practice until the flowers have matured. Biomass is harvested dried and treated to convert cannabidiolic acid (CBDA) to cannabidiol (CBD). CBD is extracted from the treated biomass using subcritical liquid carbon dioxide as solvent. The CBD is further purified by absorbing it onto an appropriate matrix and preferentially eluting it using subcritical liquid carbon dioxide. The purified CBD is recovered by allowing the carbon dioxide to evaporate.

12. The ACNFP considered whether the use of solvents as processing aids resulted in residues that require highlighting to risk managers. To assess the safety of the solvent residues that remain in the novel food, comparison was made to residue limits for other consumed products as detailed in Table 1.

**Table 1: Comparison of information on permitted residue levels for solvents used in the novel foods production compared to the proposed specification.**

<b>Solvent used</b>	<b>Available data on safe maximum level of consumption</b>	<b>Level in specification for the novel food</b>
	Commission Regulation (EU) No 231/2012.	
Subcritical liquid carbon dioxide	Content not less than 99 % v/v on the gaseous basis Regulation - 231/2012 - EN - EUR-Lex (europa.eu)	NA
	" href="#">(footnote1)	

13. The evidence presented (see Table 2 below) on composition indicates compliance with the specification for residues of solvents. When considered at the level of consumption the evidence suggests the levels of solvent residues in the novel food are below those which would represent a safety concern.

14. A HACCP statement was provided along with further details of the process and how it operates. The Committee requested further information on how the production process is effectively managed, the applicant provided reassurance on this by evidencing the control procedures are in place. The Committee concluded the production process has characterised the potential hazards and detailed the corresponding control measures sufficiently.

## **2.3 Compositional Information**

15. Results from five independent batches of the novel food demonstrated that the CBD content is produced consistently. The data is presented within Tables 2 to 6 below.

16. The data presented in Table 2 indicates CBD content is consistently above 98% purity with negligible amounts of starting materials detected across the five representative batches.

17. It is recognised that the detection and characterisation of cannabinoids in a range of food matrices is an evolving area and there are yet to be internationally recognised methods. The limitations of analytical methodology available have been subject to discussion in the Joint ACNFP and COT CBD Subgroup and remain a source of uncertainty in the assessment.

**Table 2: Compositional analysis of representative batches of cannabidiol (CBD) isolate.**

<b>Parameter</b>	<b>Method of Analysis</b>	<b>Specification/ RL</b>	<b>Batch 1</b>	<b>Batch 2</b>	<b>Batch 3</b>	<b>Batch 4</b>	<b>Batch 5</b>
CBD Purity (%)	qNMR	Corresponds to reference	100.1	98.8	99.1	98.5	98.9
CBC (mg/kg)	LC-MS/MS	2.5	2.5	2.5	2.5	2.5	2.5
CBC-A (mg/kg)	LC-MS/MS	2.5	2.5	2.5	2.5	2.5	2.5
CBD-A (mg/kg)	LC-MS/MS	2.5	15	15	15	15	15
CBD-V (mg/kg)	LC-MS/MS	2.5	> 500	> 500	> 500	> 500	> 500
CBDV-A (mg/kg)	LC-MS/MS	2.5	1	1	1	1	1
CBG (mg/kg)	LC-MS/MS	20	20	20	20	20	20

CBG-A (mg/kg)	LC-MS/MS	2.5	1.75	1.75	1.75	1.75	1.75
CBN (mg/kg)	LC-MS/MS	2.5	2.5	1.75	1.75	2.5	1.75
Delta-8-THC (mg/kg)	LC-MS/MS	2.5	15	16.8	10.7	15	2.5
Delta-9-THC (mg/kg)	LC-MS/MS	2.5	185	137	129	86	42
THC-A (mg/kg)	LC-MS/MS	5	5	5	5	5	5
THC-V (mg/kg)	LC-MS/MS	2.5	1.75	1.75	1.75	1.75	1.75
THCV-A (mg/kg)	LC-MS/MS	2.5	1	1	1	1	1

qNMR = quantitative Nuclear Magnetic Resonance Spectroscopy; RL = Reporting limit; MS/MS = tandem mass spectrometry; LC-MS = liquid chromatography-mass spectrometry.

18. Analytical data concerning the microbiological content from five independent batches of the novel food were reported (Table 3). The process in manufacturing this novel food uses subcritical liquid carbon dioxide as an extraction solvent, high temperature steps, and drying, which may mitigate the proliferation of microbes within the final product. The microbiological data presented confirm that the novel food does not raise a safety concern and consistently meets the proposed microbial specification levels.

**Table 3: The microbiological analysis of the novel food.**

Parameter*	Batch 1	Batch 2	Batch 3	Batch 4	Batch 5
------------	---------	---------	---------	---------	---------

Total viable cell count (CFU/g)	10	10	10	10	10
Total coliforms (CFU/g)	10	10	10	10	10
<i>Escherichia coli</i> ( <i>E. coli</i> ) (CFU/ g)	10	10	10	10	10
Yeasts/ Moulds (CFU/g)	100	100	100	100	100
<i>Salmonella</i> spp.(/ 25 g)	Absent	Absent	Absent	Absent	Absent

\* = Internally developed and validated methods used for analyses.

CFU = Colony forming units.

19. Results from the mycotoxin analysis for five independent representative batches of isolated CBD are presented in Table 4. The data show that the isolated CBD consistently complies with the specifications set for mycotoxins within the final product.

**Table 4: The mycotoxin analysis of the novel food.**

Parameter	Method of Analysis	Specification/ RL	Batch 1	Batch 2	Batch 3	Batch 4	Batch 5
Aflatoxin B1 (µg/kg)	HPLC-FD	0.02	0.02	0.02	0.02	0.02	0.02
Aflatoxin B2 (µg/kg)	HPLC-FD	0.02	0.02	0.02	0.02	0.02	0.02
Aflatoxin G1 (µg/kg)	HPLC-FD	0.02	0.02	0.02	0.02	0.02	0.02
Aflatoxin G2 (µg/kg)	HPLC-FD	0.02	0.02	0.02	0.02	0.02	0.02

Ochratoxin A (µg/kg)	HPLC-FD	0.02	0.02	0.02	0.02	0.02	0.02
Total aflatoxins (µg/kg)	HPLC-FD	0.08	0.08	0.08	0.08	0.08	0.08

HPLC-FD = High performance liquid chromatography - fluorescence detection.

20. Novel food products must comply with the legal requirements for heavy metal contaminants in food. Analytical data, presented for five independent batches of the novel food, demonstrated that heavy metals were present in low quantities and below established EU limits where applicable (applicable for arsenic, cadmium, mercury and lead) (Table 5).

**Table 5: Elemental Impurities in the novel food.**

<b>Parameter</b>	<b>Method of Specification/ Analysis</b>	<b>RL</b>	<b>Batch 1</b>	<b>Batch 2</b>	<b>Batch 3</b>	<b>Batch 4</b>	<b>Batch 5</b>
Elemental Impurities (Arsenic, mg/kg)	ICP-MS	0.02	0.02	0.02	0.02	0.02	0.02
Elemental Impurities (Cadmium, mg/kg)	ICP-MS	0.005	0.005	0.005	0.005	0.005	0.005
Elemental Impurities (Chromium, mg/kg)	ICP-MS	0.05	0.73	3.09	1.14	1.47	1.67



Elemental Impurities (Copper, mg/kg)	ICP-MS	0.05	0.13	0.42	0.11	0.18	0.11
Elemental Impurities (Lead, mg/kg)	ICP-MS	0.005	0.005	0.005	0.005	0.005	0.005
Elemental Impurities (Mercury, mg/kg)	ICP-MS	0.01	0.01	0.01	0.01	0.01	0.01
Elemental Impurities (Nickel, mg/kg)	ICP-MS	0.02	0.02	0.6	0.02	0.2	0.2
Elemental Impurities (Zinc, mg/kg)	ICP-MS	0.2	0.2	0.2	0.2	0.2	0.2

ICP-MS = inductively coupled plasma mass spectrometry.

21. Results from the residual solvent analysis and pesticide analysis for five independent representative batches of isolated CBD are presented in Table 6. The data show that the isolated CBD is able to consistently comply with the specifications set for residual solvents and pesticides within the final product.

**Table 6: Residual solvents and pesticides analysis in novel food.**

Parameter	Method of Analysis	Specification/ RL	Batch 1	Batch 2	Batch 3	Batch 4	Batch 5
-----------	--------------------------	----------------------	------------	------------	------------	------------	------------

Residual Solvent (Methanol, mg/kg)	GC-MS	6	6	6	6	6	6
Residual Solvent (Ethanol, mg kg)	GC-MS	6	6	6	6	6	6
Residual Solvent (Isopropyl alcohol, mg/kg)	GC-MS	6	6	6	6	6	6
Residual Solvent (Acetone, mg/kg)	GC-MS	6	6	6	6	6	6
Pesticide Screen	GC-MS; LC-MS	0.01	0.01	0.01	0.01	0.01	0.01

GC-MS = Gas chromatography-Mass Spectrometry; LC-MS = liquid chromatography-mass spectrometry.

### 2.3.1 THC as a potential contaminant in the novel food.

22. The extraction process may result in other cannabinoids remaining as contaminants. In particular, delta-9-tetrahydrocannabinol ( $\Delta$ 9-THC) and its precursor acid delta-9-tetrahydrocannabinolic acid ( $\Delta$ 9-THCA) were analysed due to the potential for toxic effects resulting from their consumption and the status of  $\Delta$ 9-THC as a controlled drug within the UK (Table 2). Along with  $\Delta$ 9-THC, other minor cannabinoids which occur at contaminant levels have the potential to play a role in the toxicity of CBD novel food products; as such, they require due consideration and monitoring to ensure the novel foods remain safe. As a result, the robustness, accuracy, and precision of the methods have been considered in interpreting the data on  $\Delta$ 9-THC and were considered appropriate in this case.

23. A literature review was undertaken as part of the assessment of CBD as a novel food, to understand the impact on the safety of foods with trace levels of contamination with  $\Delta$ 9-THC. The joint ACNFP and COT subgroup reviewed the information from literature and identified a point of departure from the European

Food Safety Authority (EFSA) opinion on  $\Delta$ 9-THC as a contaminant in milk and meat (EFSA,2015).

24. Evidence from an EFSA review by the CONTAM panel suggests a point of departure from a LOAEL (lowest observed adverse effect Level) of 0.036 mg/kg/bw/day, which is drawn from the most sensitive individuals and at the lowest dose tested in the clinical studies that were reviewed (EFSA, 2015). Uncertainty factors were then applied to identify a safe upper intake level. These included an uncertainty factor of 3 to extrapolate from a LOAEL to a NOAEL (no observed adverse effect level); which was considered appropriate considering the effects are mild to moderate in severity. A further factor of 10 was applied to account for individual variation, resulting in-total to an applied uncertainty factor of 30. This resulted in a safe upper intake level of 1  $\mu$ g /kg bw/day for  $\Delta$ 9-THC consumed as a contaminant in food (EFSA, 2015).

25. The Subgroup agreed the acute reference dose (ARfD) to be sufficiently protective to apply to the UK population. It was noted that in applying the acute reference dose EFSA has assumed that the effects seen would be the same if humans were exposed to multiple doses of THC at very low levels. The Subgroup commented that there was no data to verify this assumption, but if setting limits the dataset is the best available.

26. The analysis for delta-9-tetrahydrocannabinidiol as a potential contaminant in the novel food was declared as 42 – 185 mg/kg in the five batches tested, (Table 2), with a reporting limit of 2.5 mg/kg.

27. The levels of  $\Delta$ 9-THC, where detected in the novel food, once adjusted to the proposed use of 10 mg of CBD being consumed a day, were below the ARfD identified by EFSA of 1  $\mu$ g/kg bw/day or 70  $\mu$ g/day for a healthy adult. This level does not present a concern in terms of consumer safety for the novel food under the proposed conditions of use.

28. To ensure  $\Delta$ 9-THC levels remain consistently low in the production of CBD, THC and its precursor acid combined should be a standard substance included in the specification as relevant to all batches produced.

29. The data presented did not indicate additional hazards for inclusion in the specification.

## **2.4 Stability**

30. The stability of the novel food was assessed in real-time under the recommended storage conditions (4°C) in five batches for 6 months. Results showed that the novel food meets the specification criteria for CBD and other cannabinoid content over this time period.
31. Stability results for the CBD isolate preparation using a model system based on a 1% solution of CBD in different vegetable oils was reviewed. The accelerated shelf-life study was performed at 60°C for 24 days in clear glass packaging in full light conditions. The results of this study were interpreted and demonstrated the CBD isolate is stable for 12 months in hemp oil.
32. The data provided supports the stability of CBD isolate for a period of at least 12 months.

**2.5 Specification**

33. The applicant’s specification parameters reported in Table 7 were assessed using internationally recognised methods or determined using internally developed and validated methods. The results of the analysis are detailed in Table 2 to 6 and indicate the novel food can be produced consistently to the specification.

**Table 7: Specification of the novel food.**

Description - Cannabidiol is an amorphous powder produced by a multistep extraction process.

Parameter	Specification	Method
CBD (%)	>98.5	HPLC
CBDV (%)	1.0	HPLC
Delta-9-THC (%)	0.05	GC-MS
Terpenes (%)	2	-----

Sesquiterpenes (predominately humulene, caryophyllene and alpha bisabolol) (%)	2	GC-FID
Monoterpenes (predominantly myrcene, alpha pinene and beta pinene) (%)	1	GC-FID
<b>Other (%)</b>	1	-----
Plant cuticle waxes (predominantly long chain alkanes C27-C33) (%)	1	GC-FID
<b>Metals</b>	-----	-----
Arsenic (mg/kg)	0.02	ICPMS
Cadmium (mg/kg)	0.005	ICPMS
Chromium (mg/kg)	4.00	ICPMS
Copper (mg/kg)	1.00	ICPMS
Lead (mg/kg)	0.01	ICPMS
Mercury (mg/kg)	0.01	ICPMS
Nickel (mg/kg)	1.00	ICPMS
Zinc (mg/kg)	0.2	ICPMS
Pesticides	Compliance with national limits	GCMS-MRM / LCMS-MRM

CBD = Cannabidiol; HPLC = high-performance liquid chromatography; THC = tetrahydrocannabinol; GC-MS = Gas chromatography-mass spectrometry; GC-FID = Gas chromatography- flame ionization detection; ICPMS = Inductively coupled plasma mass spectrometry; MRM = Multiple reaction monitoring.

34. The ACNFP concluded the information provided is sufficient for the specification of CBD and appropriately characterises the novel food seeking authorisation.

## **2.6 History of Use**

35. Hemp has been widely consumed in the UK and EU as a seed oil, in tea and as an alternative to hops in beer. Extracts of hemp including CBD and synthetic CBD have not been widely consumed and are considered novel foods. While CBD products are widely available on the UK high street, indicating some consumption of CBD as a food, no applications for CBD have yet received authorisation as a novel food. Any products which are currently on the UK market therefore remain non-compliant with the novel food regulations.

36. As detailed in the COT review of the literature there has been use of both hemp derived and synthetic forms of CBD for medicinal purposes. These provide an indication of the toxicological effects that should be explored in the testing regime – primarily effects on liver, thyroid and potential impacts on reproductive organs. Also reported are behavioural effects such as somnolence (sleepiness) (COT, 2020).

37. As reported in the COT review of the publicly available data on CBD and summary data on a medicinal product, signs of adverse effects on the liver were observed at doses of CBD as low as 5 mg/kg bw/day in patients and healthy human volunteers; this dose is equivalent to 350 mg in a 70 kg adult. The data in the literature also suggested that humans might be more sensitive to the adverse effects of CBD in the liver than laboratory animals.

38. Somnolence effects were noted at doses  $\leq 10$  mg/kg bw/day in human studies. Inhibitory drug-drug interactions have also been observed with some medications when CBD is co-administered at doses of 1 mg/kg bw/day (equivalent to 70 mg in a 70 kg adult); the likelihood of effects at lower doses has not been determined. Based on the COT assessment, therefore, the FSA concluded in February 2020 that 1 mg/kg bw/day, or 70 mg in a 70 kg adult, was a pragmatic upper level of intake above which there would be clear concerns about safety.

39. It is noted that the doses used for medicinal purposes are higher than those proposed for food use. The purpose of an assessment for medicines authorisation is different to that for food and this is reflected in the data requirements. Unlike medicines, there is no risk-benefit context in foods with the requirement instead being that the products are safe. This means that outcomes that are considered an adverse event for food might not be considered as such in a medicinal study.

40. Within the literature, further human studies utilising chemically derived CBD provides further evidence of a history of synthetic CBD use (Izegelov *et al.*, 2010; Stereo Biotechs Ltd., 2020; Klotz *et al.*, 2019; Wheless *et al.*, 2019). A review by Heuestis *et al.*, 2019 of Cannabidiol Adverse effects and Toxicity notes that, while CBD is not risk-free, severe adverse events occur at doses higher than those recommended for human pharmacotherapies which are prescribed to treat forms of epilepsy. The data on previous consumption of CBD suggest areas for careful consideration in the toxicological review to understand potential effects at the lower doses used in foods.

**2.7 Proposed Use and Anticipated Intake**

41. The proposed use for the novel food is food supplements as defined byThe Food Supplements (England, Scotland and Wales) Regulations 2003 as capsules, liquid or drops in dose form.

42. The applicant initially proposed a use level of 40 mg/day CBD for the novel food in adults (excluding pregnant and lactating women, and other specifically identified vulnerable groups including those taking medication and the immunosuppressed). As a result of consultation with the applicant the proposed uses have been updated to reflect the provisional acceptable daily intake (ADI) for the use of ≥98% pure form CBD established at 10 mg per day (ACNFP and COT, 2023). The proposed maximum use levels for the novel food are outlined in Table 8.

**Table 8: Amended proposed uses and maximum use levels for the novel food.**

<b>Food Category</b>	<b>Maximum intake level from product use</b>
----------------------	--

Food Supplements (for adults) as defined in the Food Supplements (England) Regulations 2003 and other equivalent legislation in the other nations of the UK as capsules, liquid or drops in dose form intended for those 18 years of age or over. Excluding pregnant and lactating women and other specifically identified vulnerable groups. 10 mg per day

43. It is noted that consumers may be exposed to CBD from a range of food categories. The standard methodology for calculating exposure for a novel food would explore intake from a range of sources and ensure that exposure via the proposed uses would not exceed any safety level identified when consumption of the food category was analysed. It is noted that for CBD that there are already many products available. As such, the assessment has been made on the basis of identification of a maximum level of CBD that can be consumed per day. As such proposed uses will only be considered safe within the assessment when at a maximum consumption of 10mg of CBD per day from all sources (as concluded in section 2.10 of this assessment).

44. Concerns were raised by the Committee regarding the potential for foreseeable misuse of CBD if consumed in multiple formats on a single day. This is because of the increased risk of consuming CBD above the provisional acceptable daily intake (ADI). The scope of the assessment is restricted to the uses proposed and any further uses or additional food categories would be subject to the change in conditions of use process.

45. Risk managers must consider whether consumers would benefit from information on the CBD content of foods in order to ensure their consumption does not exceed the maximum intake of 10mg per day for a healthy adult.

46. As recommended in the ACNFP and COT statement on CBD of 98% purity, "The provisional ADI is recommended, subject to the existing advice to consumers that pregnant and breastfeeding women and people taking any prescription medication should avoid the consumption of CBD if possible. Consumers on regular medications should seek advice from a medical professional before using any type of CBD food product. In addition, children and prospective parents trying for a baby are advised against consumption of CBD, as are those who are immunosuppressed, due to remaining data gaps and residual uncertainties concerning the safety of CBD for these groups of consumers" (ACNFP and COT, 2023).



47. The ACNFP explored the potential for foreseeable misuse of the novel food. It was noted that the availability of multiple formats of the novel food could create conditions where exposure estimates are exceeded. It is highlighted to risk managers that they may wish to consider whether risk management measures are needed beyond those in the food supplements regulation to ensure consumers are aware of the provisional ADI of 10 mg CBD/day for the product, a dose at which it is considered that no adverse effects would be expected.

48. It is also strongly recommended that risk managers consider how consumers can be supported to manage their intake appropriately within the safe limits identified and appreciate the nature of the potential risks at higher doses, for uses that are not in dosed forms.

49. The food supplement products are to be labelled in accordance with the labelling requirements of Food Supplements (England) Regulations 2003 and the equivalent legislation in the nations of GB. The ACNFP recommended that the applicants proposed warning labelling be updated to include information on not exceeding the safe limit of 10 mg/day for a 70 kg healthy adult, that the product is not suitable for use under the age of 18 or for use during pregnancy or breastfeeding. As well as information on its suitability if the consumer is taking medication or have existing health conditions.

## **2.8 Absorption, Distribution, Metabolism and Excretion (ADME)**

50. The Absorption, Distribution, Metabolism and Excretion (ADME) of CBD are known to be complicated by the food matrix in which the CBD is delivered and are currently still being defined by professional bodies.

51. The oral bioavailability of CBD is low, indicating that it is not absorbed to any notable extent following ingestion (Mechoulam *et al.*, 2002). Published works report the bioavailability of CBD to be between 13 and 19% (Grotenhermen (2003)) or 6% (Hawksworth and McArdle (2004)). The low systemic availability was demonstrated by Martin-Santos *et al.*, 2012 and further supported by a literature search which identified the pharmacokinetics of CBD in humans (Miller *et al.*, 2018) The COT statement on CBD of 2020 noted that although CBD has low fasting bioavailability (10%), consumption with food could increase CBD uptake, by for example, 5-fold if eaten with a high fat meal. As such the potential for matrix effects that impact bioavailability cannot be ruled out. The applicant has submitted additional information in the form of a pharmacokinetic study, which was reviewed by the ACNFP (Johnson *et al.*, 2022).

52. Following oral consumption, CBD is extensively metabolised in the liver. This rapid first pass metabolism contributes to the low oral bioavailability reported in the literature (Taylor *et al.*, 2018; WHO, 2018). In vitro studies indicate that CYP3A4 and CYP2C19 are the primary hepatic enzymes responsible for first-pass metabolism of CBD; however, several other hepatic cytochrome P450 isoforms (CYP1A1, CYP1A2, CYP2C9, CYP2D6, and CYP3A5) have also demonstrated a capability of metabolising CBD (Jiang *et al.*, 2011; Zendulka *et al.*, 2016).

53. The metabolism of CBD is thought to follow two separate pathways. One is P450-mediated, in which CBD is metabolised into its major metabolite 7-COOH-CBD. This is followed by further metabolic reactions which yield the minor metabolites of CBD, including 6-OH-CBD (Devinsky *et al.*, 2018; Taylor *et al.*, 2018). The other involves decarboxylation (Kraemer *et al.*, 2019). The resultant metabolites are predominantly excreted in faeces and urine (Hawthornth and McArdle, 2004; WHO, 2018).

54. Multiple dosing with CBD is associated with a steady state concentration up to 2-fold accumulation of CBD in plasma when compared with a single dose (Taylor *et al.*, 2018). Minimal evidence of plasma accumulation has also been reported in dosing studies over 5–9 days (Millar *et al.*, 2018; Sellers *et al.*, 2013; Stott *et al.*, 2013).

55. The pharmacokinetics of CBD have been systematically reviewed by Millar *et al.*, (2018) in 24 studies, most of which assessed the administration of CBD at doses of 5–20 mg/day. This correlates to a low dose application similar to this CBD novel food application. Following oral administration, single doses of 5.4 and 10 mg CBD achieved peak serum concentrations (C<sub>max</sub>) of 0.9 and 2.5 ng/mL. The time to maximum concentration (T<sub>max</sub>) was approximately 1 hour, with a half-life between 1 and 3 hours. Given the intended use of this CBD as a food supplement, with an approximate half-life of 1 to 3 hours, with a total clearance of 6 hours, there are no significant concerns of accumulation (Millar *et al.*, 2018).

56. The ADME data provides context for interpreting the toxicological data. It is noted that the bioavailability of CBD is typically low but can be affected by food matrix. The food context for CBD could impact on CBD bioavailability. It was noted that the potential for CBD to accumulate in the body has not been examined based on the data supplied. This has been taken into account in considering the assessment factors to account for uncertainty in setting the provisional ADI.

## **2.9 Nutritional information**

57. The ACNFP sought clarification of the potential for the presence of antinutritional factors from the preparation. It was noted that hemp can contain a range of substances that could impact the digestion and absorption of nutrients from the diet. These include phytic acid (which can negatively affect the bioavailability of dietary and endogenous minerals and proteins), tannins (which can interrupt the absorption of iron), trypsin Inhibitors (which can affect protein digestion), and saponins (which at larger quantities cause gastric irritation and increase the permeability of the intestine).

58. The product is highly purified as indicated in the information on the composition. There are no substances present that would be expected to impact the digestion or absorption of nutrients from the diet.

59. The data on nutritional composition confirm that CBD has no caloric or nutritional value. The application is not intending that CBD replace another food in the diet. Consumption of the novel food at the proposed use levels is not expected to be nutritionally disadvantageous for consumers.

## **2.10 Toxicological information**

60. Toxicological studies on CBD were performed by the applicant to support the safety assessment of the novel food. The respective study reports are unpublished and claimed as confidential and proprietary data. They were considered essential in the assessment of the safety of the novel food and were reviewed by the ACNFP. How data on systemic toxicity was managed and interpreted in the context of the provisional ADI is explained in the subchronic toxicology section below (section 2.10.2).

### **2.10.1 Genotoxicity**

61. *In vitro* genotoxicity testing of CBD was conducted under Good Laboratory Practice (GLP) conditions and utilised the following OECD guidelines: *in vitro* bacterial reverse mutation test (OECD TG 471) and *in vitro* mammalian cell micronucleus test (OECD TG 487). This approach is recommended by the UK Committee on Mutagenicity and is also the basis of guidance on the preparation and submission of an application for authorisation of a novel food in the context of assimilated Regulation (EU) 2015/2283.

62. The *in vitro* bacterial reverse mutation test demonstrated that this CBD ingredient was non-mutagenic, in the absence or presence of metabolic activation. In addition, the *in vitro* mammalian cell micronucleus test demonstrated that CBD was non-clastogenic and non-aneugenic in the absence

and presence of metabolic activation.

63. The results from these *in vitro* studies support the conclusion that the novel food (>98% pure CBD) is not genotoxic. This is consistent with the view of the Committee on Mutagenicity in reviewing CBD generically as a substance from evidence available in the public domain (Committee on Mutagenicity; MUT/MIN/2020/1, 2020).

### **2.10.2 Sub-chronic Toxicity**

64. A Joint Subgroup of the ACNFP and COT was formed to address a series of questions in relation to the safety of CBD, cannabinoids and hemp-derived ingredients.

65. The applicant provided a Repeated Dose 13-week Oral Toxicity Study in Rodents [(Labcorp, study number 8462626)], which was conducted under GLP conditions and to OECD Test Guideline 408. In this 90-day study, male and female rats were dosed once a day with 0 (corn oil), 15, 30, or 60 mg/kg bw/day CBD by oral gavage at a dose volume of 5 mL/kg bw. The control group (0 mg/kg bw/day) and high-dose group (60 mg/kg bw/day) both contained 15 male and 15 female rats, and dose groups 15 and 30 mg/kg bw/day both contained 10 male and 10 female rats. Of the control group and high-dose group, 5 males and 5 females were allocated to the recovery period and retained not dosed for 4-weeks.

66. The applicant concluded a NOAEL of 60 mg/kg bw/day, the subgroup reviewed the data and concluded a NOAEL from the study of 30 mg/kg bw/day, based on liver enlargement and histopathological changes in the adrenal. The subgroup considered the critical effect on the liver to be the NOAEL and thus a point of departure (POD) of 30 mg/kg was confirmed by the subgroup.

67. The Subgroup reviewed the data and concluded the effects seen at 30 mg/kg bw/day were minimal but not adverse, however the effects seen at 60 mg/kg bw/day are potentially adverse. As such the subgroup considered that the NOAEL for the study be 30 mg/kgbw/day. Review of the study by the subgroup supported the conclusion that it was of sufficient quality to support the safety of the novel food.

68. In addition to the data submitted by the applicant there is a body of evidence on the effect of 98% or greater CBD. In order to take account of all pertinent data and to put the individual assessment in the context of the totality of relevant evidence for the active substance. The data from this application was compared to the wider body of evidence.

69. A weight of evidence approach allowed the Subgroup to identify a provisional ADI for CBD ingredients of >98% purity of 0.15 mg/kg bw/day or 10 mg per day for a 70 kg healthy adult (Joint position paper from the ACNFP and COT; FSA consumer advice published in October 2023). This value was identified to be protective of the most sensitive known effects in the liver and thyroid parameters and included consideration of data gaps and uncertainties.

70. The dataset includes several studies where highly purified CBD has been tested. Given the low level of contaminants, it is reasonable to consider that these represent the effect of CBD as a substance and are therefore relevant to other novel foods with similar compositions.

71. It was considered whether the wider data and therefore the provisional ADI for CBD of 98% or greater purity was relevant to the review of this novel food. It was considered appropriate, on the basis that the test substance used in the study to support the novel food was 98% pure and the compositional data was consistent with a highly purified CBD. The contaminants present were not suggestive of a significant impact on the toxicology. The point of departure in the form of a NOAEL from the study submitted to support safety of this novel food once corrected for CBD content, is consistent with the range of the points of departure used to develop the provisional ADI (ACNFP and COT, 2023). The NOAEL was also based on the same effect - impacts on the liver. The uncertainty factors identified in the provisional ADI would also apply to the applicant's submitted study (Labcorp, study number 8462626 unpublished) for the same reasons as identified in the provisional ADI statement. It was, therefore, considered scientifically appropriate to apply the provisional ADI of 0.15 mg/kg bw/day or 10 mg/day as identified in the joint statement of the ACNFP and COT on  $\geq 98\%$  pure forms of CBD to the novel food in this application.

## **2.11 Allergenicity**

72. This CBD isolate comprises >98% CBD and the production process for CBD does not introduce any risk of allergenic potential. As a chemical entity the potential for IgE mediated food allergy is unlikely.

73. Given CBD as a substance is not considered allergenic, the allergenicity assessment considered whether the other 2% of the novel food's composition was likely to be allergenic or elicit food allergic reactions. It was noted that none of the raw materials or processing aids used in the production process are derived from or contain any of the allergenic food ingredients specified under Regulation (EU) No 1169/2011 on the provision of food information to consumers. This suggests

the potential to elicit reactions in those sensitive to those foods is unlikely.

74. The novel food is unlikely to trigger allergic reactions in the target population under the proposed conditions of use.

### **3. Discussion**

75. The novel food is a CBD isolate ingredient from industrial hemp containing >98% CBD, produced using a multi-step manufacturing process.

76. This CBD isolate is intended to be used as a food ingredient in food supplements for adults excluding pregnant and lactating women and other specifically identified vulnerable groups at a defined intake for each product type of up to 10 mg CBD per day; it is not intended to replace any food.

77. In October 2023, the Joint ACNFP and COT subgroup identified a provisional acceptable daily intake (ADI) of 10 mg per day (0.15 mg/kg bw/day) for CBD products containing 98% CBD or above, such as the novel food discussed in this assessment. A weight of evidence approach was used to arrive at a provisional ADI of 10 mg/day (0.15 mg/kg bw/day). The most sensitive human health effects, which this provisional ADI protects against, are seen consistently in the liver and thyroid in a number of studies using  $\geq 98\%$  pure CBD. This value also takes account of the lack of human-based long-term evidence or evidence regarding potentially vulnerable groups.

78. Based upon the dossier of evidence provided by the applicant, the safety of the novel food was reviewed and evidence to reach a conclusion on safety provided. The evidence presented by the applicant was then compared to the wider data set on CBD and the evidence presented is consistent with evidence presented to support the development of a provisional ADI of 10 mg/day for CBD of 98% purity or above. As such it is appropriate to apply the provisional ADI to this novel food.

79. This is subject to the existing advice to consumers that pregnant and breastfeeding women and people taking any prescription medication should avoid the consumption of CBD. Consumers on regular medications should seek advice from a medical professional before using any type of CBD food product. In addition, children and prospective parents trying for a baby are advised against consumption of CBD, as are those who are immunosuppressed, due to remaining data gaps and residual uncertainties concerning the safety of CBD for these groups of consumers. These contraindications would also apply to this novel food.

80. The maximum safe exposure for healthy adults of 70 kg as identified in the provisional ADI is 10 mg per day from all food sources. If the inclusion level of this CBD isolate leads to an intake per individual serving of each product type of 10 mg/day, multiple intakes of food products containing CBD on the same day should be avoided to support minimising exposure to below the provisional ADI.

## 4. Conclusions

81. The ACNFP has undertaken a review of this CBD isolate and concluded that the novel food is safe under the proposed conditions of use and does not pose a safety risk to human health. The proposed uses are not considered nutritionally disadvantageous.

82. These conclusions were supported by the information in the novel food dossier submitted by the applicant plus the supplementary information and could not have been reached without the following data claimed as proprietary by the applicant:

- annexes to the dossier which relate to the identity of the novel food, the production process, stability, methods of analysis, particle size analysis and toxicology.
- *in vitro* bacterial reverse mutation test [(Labcorp study number 8462622)], *in vitro* mammalian cell micronucleus test [(Labcorp study number 8242623)] and 13-weeks repeat dose feeding study [(Labcorp study number 8462626)].

83. The members of the ACNFP during the course of the assessment who were; Dr Camilla Alexander White, Dr Anton Alldrick, Dr Kimon Andreas Karatzas, Alison Austin, Professor George Bassel, Dr Mark Berry, Dr Christine Bosch, Professor Dimitris Charalampopoulos, Dr Catharina Edwards, Professor Susan Fairweather-Tait, Professor Paul Frazer, Dr Hamid Ghoddusi, Professor Andy Greenfield, Professor Wendy Harwood, Professor Huw D. Jones, Dr Ray Kemp, Dr Elizabeth Lund, Professor Harry J. McArdle, Mrs Rebecca McKenzie, Professor Clare Mills, Dr Antonio Peña-Fernández, Dr Lesley Stanley, Professor Hans Verhagen, Dr Maureen Wakefield, Dr Meera Cush, Dr Sophie Foley, Dr Lynn McIntyre, Dr Isabel Skypala and Professor Bruce Whitelaw.

84. To note, interests were received from members of the ACNFP, Dr Alldrick declared a potential interest relating to his previous employment and this was considered a potential conflict and as a result he was not present for discussions of CBD by the Committee. Emeritus Prof Harry McArdle declared an interest from his work with EFSA's Nutritional Novel Food and Food Allergens

Panel in considering data requirements for CBD. While not seen as a conflict, to avoid Prof McArdle being subject to information that would influence his EFSA work, it was agreed that he would not be present in discussions for CBD by the ACNFP but could supply comments for consideration by the Committee upon review of the minutes.

## 5. References

Advisory Committee on Novel Foods and Processes (ACNFP) and Committee on Toxicity (COT), 2023. Joint position paper from the Advisory Committee on Novel Foods and Processes (ACNFP) & Committee on Toxicity (COT) on establishing a provisional acceptable daily intake (ADI) for pure form ( $\geq 98\%$ ) cannabidiol (CBD) in foods, based on new evidence. Published October 2023 DOI:

<https://doi.org/10.46756/sci.fsa.zcg392>.

AOAC (2007) Official Methods of Analysis. 18th Edition, Association of Official Analytical chemists, Gaithersburg.

AOAC Cannabis Analytical Science Program (CASP). "AOAC SMPR® 2019.002." Standard Method Performance Requirements (SMPRs®) for Identification and Quantitation of Selected Residual Solvents in Cannabis-Derived Materials, 9 Oct. 2019.

AOAC Official Method 2018.10 Determination of Cannabinoids in Cannabis sativa Dried Flowers and Oils Liquid Chromatography with UV Detection First Action 2018.

Audino, Susan *et al.* "AOAC SMPR® 2017.001. Standard Method Performance Requirements (SMPRs) for Quantitation of Cannabinoids in Cannabis Concentrates." *Journal of AOAC International* 100 4 (2017): 1200-1203.

Borman, Phillip, and David Elder. "Q2(R1) Validation of Analytical PROCEDURES." ICH Quality Guidelines, 2017, pp. 127-166., doi: 10.1002/9781118971147.ch5 <https://doi.org/10.1002/9781118971147.ch5>.

Collins, Arianna C. Extraction Procedures HPLC (AC001). 2020. Midwest Extraction Services Analytical Laboratory, Minnesota.

Committee On Mutagenicity Of Chemicals In Food, Consumer Products And The Environment, (MUT/MIN/2020/1), 2020; [Committee on Mutagenicity minutes from meeting on 20th February 2020](#).



Committee On Toxicity Of Chemicals In Food, Consumer Products And The Environment, (TOX/2020/02), 2020; [Committee on Toxicity CBD Update](#).

Crujisen, Hans & Poitevin, Eric & Brunelle, Sharon & Almeida, S & Braun, U & Connelly, M & Giuliani, L & Huertas, Raquel & Hui, S & Ikeuchi, Y & Jaudzems, G & Kimura, S & Kittleson, J & Larkin, G & Li, F & McMahon, A & Nagatoshi, M & Piccon, I & Postma, M & Xindong, G. (2019). Determination of Minerals and Trace Elements in Milk, Milk Products, Infant Formula, and Adult Nutrition: Collaborative Study 2011.14 Method Modification. Journal of AOAC INTERNATIONAL. 102. 1845-1863. 10.1093/jaoac/102.6.1845. <https://doi.org/10.5740/jaoacint.19-0073>.

Determination of Chromium, Selenium, and Molybdenum in Infant Formula and Adult Nutritional Products - Inductively Coupled Plasma-Mass Spectrometry: Collaborative Study, Final Action 2011.19. J AOAC Int. 2015 Jun 24. doi: 10.5740/jaoacint.15139. Epub ahead of print. PMID: 26111082. <https://doi.org/10.5740/jaoacint.15139>.

Devinsky O, Patel AD, Thiele EA, Wong MH, Appleton R, Harden CL, Greenwood S, Morrison G and Sommerville K, 2018. Randomized, dose-ranging safety trial of cannabidiol in Dravet syndrome [GWPCARE1 Part A Study Group]. Neurology, 90, e1204-e1211. <https://doi.org/10.1212/WNL.0000000000005254>.

EC, 2017. Commission Implementing Regulation (EU) 2017/2469 of 20 December 2017 laying down administrative and scientific requirements for applications referred to in Article 10 of Regulation (EU) 2015/2283 of the European Parliament and of the Council on novel foods. [Commission Implementing Regulation \(EU\) 2017/2469](#).

EFSA CONTAM Panel (Panel on Contaminants in the Food Chain), 2015. Scientific opinion on the risks for human health related to the presence of Tetrahydrocannabinidiol (THC) in milk and other food from animal origin. DOI, [10.2903/j.efsa.2015.4141](https://doi.org/10.2903/j.efsa.2015.4141).

EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2021. Guidance on the preparation and submission of an application for authorisation of a novel food in the context of Regulation (EU) 2015/2283 (Revision 1). Published 26 March 2021. <https://doi.org/10.2903/j.efsa.2011.2170>.

EU (2015) Regulation (EU) 2015/2283 of the European Parliament and of the Council of 25 November 2015 on novel foods, amending Regulation (EU) No 1169/2011 of the European Parliament and of the Council and repealing Regulation (EC) No 258/97 of the European Parliament and of the Council and

Commission Regulation (EC) No 1852/2001 [Regulation \(EU\) 2015/2283 of the European Parliament and of the Council](#).

Grotenhermen, F. (2003) 'Pharmacokinetics and pharmacodynamics of cannabinoids', *Clinical Pharmacokinetics*, 42(4), 327–360.  
<https://doi.org/10.2165/00003088-200342040-00003>.

Hawksworth, G. and McArdle, K. (2004) 'Metabolism and pharmacokinetics of cannabinoids', in Guy, G, Whittle, B and Robson, P (eds.), *The Medicinal Uses of Cannabis and Cannabinoids*. London: London Pharmaceutical Press. pp. 205–228. Google Scholar.

Huestis MA, Solimini R, Pichini S, Pacifici R, Carlier J, Busardò FP. Cannabidiol Adverse Effects and Toxicity. *Curr Neuropharmacol*. 2019;17(10):974-989. doi: [10.2174/1570159X17666190603171901](https://doi.org/10.2174/1570159X17666190603171901).

ICH Harmonised Tripartite Guideline, Validation of Analytical Procedures: Text and Methodology Q2(R1) [ICH\\_Q2\(R2\)\\_Guideline\\_2023\\_1130.pdf](#).

ICH Q3C(R6) Maintenance on the Guideline for Residual Solvents. [Q3C-R6\\_Guideline\\_ErrorCorrection\\_2019\\_0410\\_0.pdf \(ich.org\)](#).

Izegelov D, Davidson E, Barasch D, Regev A, Domb AJ, Hoffman A. Pharmacokinetic investigation of synthetic cannabidiol oral formulations in healthy volunteers. *Eur J Pharm Biopharm*. 2020 Sep;154:108-115. doi: [10.1016/j.ejpb.2020.06.021](https://doi.org/10.1016/j.ejpb.2020.06.021).

Jiang R, Yamaori S, Takeda S, Yamamoto I and Watanabe K, 2011. 'Identification of cytochrome P450 enzymes responsible for metabolism of cannabidiol by human liver microsomes', *Life Sciences*, 89(5–6), 165–170.  
<https://doi.org/10.1016/j.lfs.2011.05.018>.

Johnson D.A, Funnell M, Heaney L, Cable T, Bailey S, Clifford T and James L.J. A Pharmacokinetic comparison of 70 mg cannabidiol ingested orally via gelatin capsules versus sublingual drops. Loughborough University. January 2022 [unpublished].

Joint position paper from the Advisory Committee on Novel Foods and Processes (ANCFP) & Committee on Toxicity (COT) on establishing a provisional acceptable daily intake (ADI) for pure form ( $\geq 98\%$ ) cannabidiol (CBD) in foods, based on new evidence; October 2023. <https://doi.org/10.46756/sci.fsa.zcg392>.

Kraemer M, Broecker S, Madea B, Hess C. Decarbonylation: A metabolic pathway of cannabidiol in humans. *Drug Test Anal.* 2019 Jul;11(7):957-967. doi: 10.1002/dta.2572. Epub 2019 Mar 20. PMID: 30698361.  
<https://doi.org/10.1002/dta.2572>.

Klotz KA, Grob D, Hirsch M, Metternich B, Schulze-Bonhage A, Jacobs J. Efficacy and Tolerance of Synthetic Cannabidiol for Treatment of Drug Resistant Epilepsy. *Front Neurol.* 2019 Dec 10;10:1313. DOI: [10.3389/fneur.2019.01313](https://doi.org/10.3389/fneur.2019.01313).

Labcorp, 2022 [unpublished]. Dated: 06 April 2022. Labcorp Early Development Laboratories Limited, Huntingdon, Cambridgeshire, UK. Study Title: Cannabidiol: bacterial reverse mutation test. Final report. Confidential. (Study No: 8462622).

Labcorp, 2022 [unpublished]. Dated: 06 April 2022. Labcorp Early Development Laboratories Limited, Huntingdon, Cambridgeshire, UK. Study Title: Cannabidiol: In vitro micronucleus test in human lymphocytes. Final report. Confidential. (Study No: 8462623).

Labcorp, 2022 [unpublished]. Dated: 28 July 2022. Labcorp Early Development Laboratories Limited, Huntingdon, Cambridgeshire, UK. Study Title: Cannabidiol: Toxicity study by oral gavage administration to Han Wistar Rats for 13 weeks followed by a 4-week recovery period. Final report. Confidential. (Study No: 8462626).

Lehotay, Steven J. "Supplemental Information for: Determination of Pesticide Residues in Foods by Acetonitrile Extraction and Partitioning with Magnesium Sulfate: Collaborative Study." *Journal of AOAC INTERNATIONAL*, vol. 90, no. 2, 2007. <https://doi.org/10.1093/jaoac/90.2.1SUP>.

Mandrioli, Mara, *et al.* "Fast Detection of 10 Cannabinoids by RP-HPLC-UV Method in Cannabis Sativa L." *Molecules*, vol. 24, no. 11, 2019, p. 2113.  
<https://doi.org/10.3390/molecules24112113>.

Martin-Santos, R., Crippa, J. A., Batalla, A., Bhattacharyya, S., Atakan, Z., Borgwardt, S., Allen, P., Seal, M., Langohr, K., Farré, M., Zuardi, A. W and McGuire, P. K. (2012). Acute effects of a single, oral dose of d9-tetrahydrocannabinol (THC) and cannabidiol (CBD) administration in healthy volunteers. *Current pharmaceutical design*, 18(32), 4966-4979.  
<https://doi.org/10.2174/138161212802884780>.

'MDA', Misuse of Drugs Act 1971, UK Public General Acts 1971 c. 38,  
<https://www.legislation.gov.uk/ukpga/1971/38/contents>.

Mechoulam R, Parker LA and Gallily R, 2002. Cannabidiol: an overview of some pharmacological aspects. *Journal of Clinical Pharmacology*, 42, 115-195.  
<https://doi.org/10.1002/j.1552-4604.2002.tb05998.x>.

Millar SA, Stone NL, Yates AS and O'Sullivan SE, 2018. A Systematic Review on the Pharmacokinetics of Cannabidiol in Humans. *Frontiers in Pharmacology*, 9, 1365 [13pp, plus supplementary table]. <https://doi.org/10.3389/fphar.2018.01365>.

OECD (Organisation for Economic Co-operation and Development), 1997. Bacterial reverse mutation test. In OECD guidelines for the testing of chemicals. OECD guideline No 471 (updated & adopted: 21 July 1997). Paris, France: Organisation for Economic Co-operation and Development (OECD).  
<https://doi.org/10.1787/9789264071247-en>.

OECD (Organisation for Economic Co-operation and Development), 1998. OECD principles of good laboratory practice. Series on principles of good laboratory practice and compliance monitoring, No. 1 (ENV/MC/CHEM(98) 17). Paris, France: Organisation for Economic Co-operation and Development (OECD), Environment Directorate, Chemicals Group and Management Committee.  
<https://doi.org/10.1787/9789264078536-en>.

OECD (Organisation for Economic Co-operation and Development), 2016. In vitro mammalian cell micronucleus test. In OECD guidelines for the testing of chemicals. OECD guideline No 487 (updated & adopted: 29 July 2016). Paris, France: Organisation for Economic Cooperation and Development (OECD).  
<https://doi.org/10.1787/9789264264861-en>.

OECD (Organisation for Economic Co-operation and Development), 2018. Repeated dose 90-day oral toxicity study in rodents. In OECD guidelines for the testing of chemicals. OECD guideline No 408 (updated and adopted 27 June 2018). Paris, France: Organisation for Economic Cooperation and Development (OECD). <https://doi.org/10.1787/9789264070707-en>.

[Regulation \(EU\) No 1169/2011 of the European Parliament and of the Council of 25 October 2011 on the provision of food information to consumers, amending Regulations \(EC\) No 1924/2006 and \(EC\) No 1925/2006 of the European Parliament and of the Council, and repealing Commission Directive 87/250/EEC, Council Directive 90/496/EEC, Commission Directive 1999/10/EC, Directive 2000/13/EC of the European Parliament and of the Council, Commission Directives 2002/67/EC and 2008/5/EC and Commission Regulation \(EC\) No 608/2004 \(Text with EEA relevance\) \(legislation.gov.uk\)](#).

Sellers EM, Schoedel K, Bartlett C, Romach M, Russo EB, Stott CG, Wright S, White L, Duncombe P, Chen CF. A Multiple-Dose, Randomized, Double-Blind, Placebo-Controlled, Parallel-Group QT/QTc Study to Evaluate the Electrophysiologic Effects of THC/CBD Spray. Clin Pharmacol Drug Dev. 2013 Jul;2(3):285-94.

<https://doi.org/10.1002/cpdd.36>.

Stott CG, White L, Wright S, Wilbraham D, Guy GW. A phase I study to assess the single and multiple dose pharmacokinetics of THC/CBD oromucosal spray. Eur J Clin Pharmacol. 2013 May;69(5):1135- 47. <https://doi.org/10.1007/s00228-012-1441-0>.

Stero Biotech Ltd. Phase 2a Study to Evaluate the Safety, Tolerability and Efficacy of Cannabidiol as a Steroid-sparing Therapy in Steroid-dependent Crohn's Disease Patients. ClinicalTrials.gov Identifier: NCT04056442. Last Update Posted : June 30, 2020. Available online at: [A Phase 2a Study to Evaluate the Safety, Tolerability and Efficacy of Cannabidiol as a Steroid-sparing Therapy in Steroid-dependent Crohn's Disease Patients](#).

Labcorp, 2021 [unpublished]. Dated: 28 July 2022. Labcorp Early Development Laboratories Limited, Huntingdon, Cambridgeshire, UK. Study Title: Cannabidiol: Toxicity study by oral gavage administration to Han Wistar Rats for 13 weeks followed by a 4-week recovery period. Final report. Confidential. (Study No: 8462626).

The Food Supplements (England) Regulations 2003, No. 1387, England; [The Food Supplements \(England\) Regulations 2003](#).

[The Food Supplements \(Wales\) Regulations 2003 \(legislation.gov.uk\)](#).

The Food Information Regulations 2014, No.1855; [The Food Information Regulations 2014](#).

Taylor L, Gidal B, Blakey G, Tayo B, Morrison G. A Phase I, Randomized, Double-Blind, Placebo Controlled, Single Ascending Dose, Multiple Dose, and Food Effect Trial of the Safety, Tolerability and Pharmacokinetics of Highly Purified Cannabidiol in Healthy Subjects. CNS Drugs. 2018 Nov;32(11):1053-1067. <https://doi.org/10.1007/s40263-018-0578-5>.

Wheless JW, Dlugos D, Miller I, Oh DA, Parikh N, Phillips S, Renfro JB, Roberts CM, Saeed I, Sparagana SP, Yu J, Cilio MR; INS011-14-029 Study Investigators. Pharmacokinetics and Tolerability of Multiple Doses of Pharmaceutical-Grade Synthetic Cannabidiol in Pediatric Patients with Treatment-Resistant Epilepsy. CNS Drugs. 2019 Jun;33(6):593-604. doi: [10.1007/s40263-019-00624-4](https://doi.org/10.1007/s40263-019-00624-4).

WHO (World Health Organization), 2018. Cannabidiol (CBD) critical review report. 40th Meeting - Expert Committee on Drug Dependence, Jun. 4-7, 2018, Geneva. Available online: [Cannabidiol \(CBD\) critical review report](#).

Zendulka, O., Dovrtělová, G., Nosková, K., Turjap, M., Šulcová, A., Hanuš, L and Juřica, J. (2016). Cannabinoids and Cytochrome P450 Interactions. *Current drug metabolism*, 17(3), 206–226.

<https://doi.org/10.2174/1389200217666151210142051>.

## Abbreviations

ACNFP Advisory Committee on Novel Foods and Processes

ADI Acceptable Daily Intake

ADME Absorption, Distribution, Metabolism and Excretion

AOAC Association of Official Analytical Chemists

ARfD Acute Reference Dose

aw Water activity

bw Body weight

CAS Chemical Abstracts Service

CBD Cannabidiol

Cmax Peak serum concentration

COT Committee on Toxicity

CFU Colony Forming Unit

DSC    Differential scanning calorimetry

EC     European Commission

EDA    Energy dispersive X-ray

EFSA   European Food Safety Agency

EMA    Environmental Medicines Agency

EU     European Union

FDA    Food and Drug Administration (USA)

FSA    Food Standards Agency

FSS    Food Standards Scotland

FTIR   Fourier-Transform Infrared Spectroscopy

GC     Gas Chromatography

GLP    Good Laboratory Practice

HACCP Hazards Analysis and Critical Control Points

HPLC   High Performance Liquid Chromatography

HS     Headspace

ICP-MS Inductively coupled plasma mass spectrometry

LC-MS Liquid chromatography-mass spectrometry

LOAEL Lowest Observable Adverse Effect Level

LOD Limit of Detection

LOQ Limit of quantification

MS Mass Spectrometry

ND Not Determined

NOEL No Observed Effect Level

NM Not measured

OECD Organisation for Economic Co-operation and Development

Ph.Eur European Pharmacopeia

TGA Thermogravimetric analysis

Tmax Time to maximum concentration

USP United States Pharmacopeia

UV Ultra-violet

XRD X-ray diffraction



1.