

Meeting

6'-Sialyllactose Sodium Salt Discussion Paper

Committee Paper for Discussion - ACNFP/162/11

Advisory Committee For Novel Foods and Processes.

**Application for Authorisation as a Novel Food for 6'-
Sialyllactose Sodium Salt, from Kyowa Hakko Bio Co. Ltd**

Application number RP1478

Issue

1. An application has been received under the novel food authorisation process (regulation 2015/2283) for 6'-Sialyllactose sodium salt (6'-SL), a Human Milk Oligosaccharide (HMO).
2. The Committee is asked to advise whether the available data provides an adequate basis for a risk assessment, and whether the novel food is safe and not nutritionally disadvantageous under the proposed use and use levels.

Background

3. On the 10th March 2022 the FSA and FSS received the submission for 6'-SL from Kyowa Hakko Bio Co., Ltd.
4. The novel food ingredient is produced via fermentation using a genetically modified strain of E. coli W followed by a series of purification steps involving isolation, concentration, and spray-drying. The applicant proposes to use the novel food in conventional foods (dairy products and analogues and cereal bars), beverages, foods for specific groups (including Foods for Special Medical Purposes, total diet replacement for weight control, infant formulas, follow-on formula, foods for infants and young children) and in food supplements as an

alternative source of 6'-SL in the absence of other exposure sources.

5. Since the submission of the present application, an authorisation for 6'-SL as a novel food on the GB market was granted in June 2022 for another applicant who sought and received data protection. In accordance with Article 26 of Regulation 2015/2283 a subsequent applicant can obtain authorisation for 6'-SL as a NF on the condition that no reference is made to the protected proprietary scientific evidence or data.

6. The present application is submitted as a full application with newly provided scientific data and evidence to support an authorisation. At EU level, an opinion on the safety for 6'-SL as a NF as submitted by Kyowa Hakko Bio Co., Ltd, was adopted by EFSA in April 2023.

7. The application is seeking an authorisation for the same uses and use levels as the previous application for 6'-SL authorised in 2022. As this is a separate applicant there are differences in the strain of microbe used, their internal production, systems and the testing used as this was a proprietary method.

8. The application dossier is attached as Annex A. The request for further information letters (with responses) are attached as Annex B. All supporting documents for all sections of the application, which have been updated following the request for further information, can be found in Annex C. All annexes contain confidential information.

9. As part of a trial for a new way of working for some applications, a draft Committee Advice Document for this application has been prepared for members consideration, alongside this introductory paper. This reflects that there have been several reviews for breast milk sugars and as such the key considerations have been discussed by the Committee in the past. Members are invited to review and comment on the draft found in Annex D.

This application

Identification

10. The novel food ingredient, 6'-SL, is a sialylated oligosaccharide composed of three molecules N acetylneuraminic acid (NeuAc), D galactose and D glucose and is identical in structure to 6'-SL found in human milk (Annex A: p12 - 14 of dossier). Purity by specification is $\geq 82\%$ (dry weight basis (dwb)) and the final novel food product includes low levels of related saccharide molecules N

acetylneuraminic acid ($\leq 9\%$ w/w), 6'-Sialyllactulose ($\leq 5\%$ w/w), D-glucose ($\leq 3\%$ w/w), D-lactose ($\leq 3\%$ w/w) and 3'-Sialyllactose sodium salt ($\leq 1\%$ w/w). Identity confirmed by LC-MS spectrometry and ^1H NMR and ^{13}C NMR spectroscopy (Annex C [Annex A]). The final composition is verified by HPLC-CAD (Annex C [Annex G]).

11. The novel food product is expected to contain 3'-Sialyllactose sodium salt (3'-SL) in low levels. An authorisation for 3'-SL as a novel food on the GB market exists but the application is held by a named applicant with use of protected proprietary data. A separate application for authorisation for 3'-SL as a novel food was submitted by Kyowa Hakko bio co., ltd on March 10th 2022.

12. The applicant has provided a range of data that characterises the identity of the genetically modified fermentation microorganism *E. coli* W, including taxonomic classification (Annex A; p 14 – 15, Table 2.a.2-1 of dossier) and information on the modification and donor organisms (Annex C [Annex B]). Identity verified by Whole Genome Sequence (WGS) and highly homologous to the *E. coli* W strain per the American Type Culture Collection (ATCC) ATCC 9637 (Annex C [Annex C]).

Production Process

13. The production method is described in detail (Annex A; p21 of dossier) and a schematic overview is provided (Annex A; p 22 of dossier and Annex C [Annex E]).

14. The production process involves a 2-step fermentation and purification process. The initial fermentation step involves culture of the production microorganism in a defined nutrient media under controlled conditions. Main fermentation involves inoculation of the cultured *E. coli* W into a nutrient media to synthesise 6'-SL in the presence of glucose and lactose.

15. The second processing step involves isolation and purification of 6'-SL using filtration and cationic and anionic exchange followed by concentration and spray-drying.

16. Additional details on pH and controls were provided by the applicant (Annex B; p 3 of response letter).

17. The acceptance criteria and specifications for the raw materials and processing aids have been provided (Annex A; p 19 – 21 and Table 2.b.1.2-1 of dossier) (Annex C [Annex D]).

18. Initially the applicant used a media containing soy peptone for the fermentation step. This element of the production process has been updated to remove the presence of soy to minimise the risk for those allergic to this food. The applicant has provided details to show whether the novel food is altered by use of the alternative media and the potential impact on the risks for the novel food. The applicant states that the additional raw materials do not introduce new hazards or changes to parameters in the production process (Annex B; p 3 of response letter).

19. The applicant has provided a HACCP Plan (Annex C [Annex E]) to include critical control measures alongside justifications. Quality control is described (Annex A; p 22 of dossier).

Composition

20. The applicant has reported analytical data for a number of independent batches of 6'-SL (Annex A; p 24 - 30 of dossier) (Annex C [Annex F]). Additional analyses were requested to provide comparable data for all parameters on batches representative of the final product and as intended to be manufactured (Annex B; p 15 - 18 and Table 3.a-3 of response letter). The data indicates that the manufacturing process results in a consistent final ingredient that meets the proposed specifications for the novel ingredient - see Tables 1-6 below. The CoA's are provided in Annex C [Annex F]. The new media has little impact on the composition of the novel ingredient and as such this was accepted as justification for providing analysis for 3 further batches with the new media

Table 1. Compositional batch analysis of 6'-SL manufactured with a fermentation media containing soy peptone: physicochemical properties, carbohydrate content and purity.

		Batch A	Batch B	Batch C	Batch D	Batch E
Parameter	Specification					
Physicochemical						

pH	4.0 – 9.0	6.4	6.5	6.5	6.5	6.5
Composition						
Purity (6'-SL)	≥82 (% dwb)	87	92	90	92	92
NeuAc	≤9 w/w %	5.1	3.5	4.9	4.3	5.4
D-glucose	≤3 w/w %	ND	ND	ND	ND	ND
D-lactose	≤3 w/w %	≤0.05	≤0.05	≤0.05	≤0.05	≤0.05
6'-sialyllactulose	≤5 w/w %	0.4	0.4	0.5	0.5	0.4
3'-sialyllactose sodium salt	≤1 w/w %	ND	ND	ND	ND	ND
Sum of other carbohydrates (2)	Not established	6.5	3.3	3.7	2.5	1.4
Water	≤10.5 w/w %	5.3	5.0	5.4	5.6	5.0
Sodium	≤5.0 w/w %	3.8	3.8	3.8	3.7	3.8
Residual protein (internal method)	≤10 mg/kg (0.001%)	≤1 (LOQ)	≤1 (LOQ)	≤1 (LOQ)	≤1 (LOQ)	≤1 (LOQ)
Residual protein (Bradford assay)	≤100 mg/kg (1)	NT	NT	≤100	NT	NT

6'-SL: 6'-sialyllactose sodium salt; NeuAc: N acetylneuraminic acid; dwb: dry weight basis; NT: not tested; ND: not detected; LOQ: limit of quantification. (1) Evaluated using a test limit at 100 ppm; (2) Sum of carbohydrates such as NeuAc, D-glucose, D-lactose, 6'-sialyllactulose and 3'-sialyllactose sodium salt

Table 2. Compositional batch analysis of 6'-SL manufactured with an updated fermentation media without soy peptone: physicochemical properties, carbohydrate content and purity.

Parameter	Specification	Batch		
		H	I	J
Physicochemical				
pH	4.0 – 9.0	6.1	5.9	5.8
Composition				
Purity (6'-SL)	≥82 (% dwb)	91	93	95
NeuAc	≤9 w/w %	1.8	0.1	0.2
D-glucose	≤3 w/w %	0.05	0.05	0.05
D-lactose	≤3 w/w %	0.05	0.05	0.05
6'-sialyllactulose	≤5 w/w %	NT	NT	NT
3'-sialyllactose sodium salt	≤1 w/w %	NT	NT	NT

Sum of other carbohydrates (2)	Not established	4.6	5.6	3.6
Water	≤10.5 w/w %	7.9	8.4	8.6
Sodium	≤5.0 w/w %	3.8	3.7	3.8
Residual protein (internal method) ≤10 mg/kg (0.001%)	NT	NT	NT	NT
Residual protein (Bradford assay) ≤100 mg/kg (1)	≤100	≤100	≤100	≤100

6'-SL: 6'-sialyllactose sodium salt; NeuAc: N acetylneuraminic acid; dwb: dry weight basis; NT: not tested; ND: not detected; LOQ: limit of quantification. (1) Evaluated using a test limit at 100 ppm; (2) Sum of carbohydrates such as NeuAc, D-glucose, D-lactose, 6'-sialyllactulose and 3'-sialyllactose sodium salt

Table 3. Compositional batch analysis of 6'-SL manufactured with a fermentation media containing soy peptone: heavy metal contaminants.

Parameter	Specification	Batch A	Batch B	Batch C	Batch D	Batch E
Heavy metals:						
Arsenic	≤0.2 mg/kg	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)
Cadmium	≤0.2 mg/kg	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)
Lead	≤0.2 mg/kg	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)

Mercury	≤0.2 mg/kg	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)	≤0.05 (LOQ)
Iron	≤10 mg/kg	0.3	0.2	0.3	0.3	0.6

LOQ: limit of quantification.

Table 4. Compositional batch analysis of 6'-SL manufactured with an updated fermentation media without soy peptone: heavy metal contaminants.

		Batch H	Batch I	Batch J
Parameter	Specification			
Heavy metals:				
Arsenic	≤0.2 mg/kg	0.01	0.01	0.03
Cadmium	≤0.2 mg/kg	0.01	0.01	0.01
Lead	≤0.2 mg/kg	0.02	0.03	0.02
Mercury	≤0.2 mg/kg	0.004	0.004	0.004
Iron	≤10 mg/kg	NT	NT	NT

NT: not tested.

Table 5. Compositional batch analysis of 6'-SL manufactured with a fermentation media containing soy peptone: microbial contaminants.

Batch A Batch B Batch C Batch D Batch E

Parameter	Specification					
Microbial parameters:						
Aerobic plate count	≤1,000 CFU/g	10	10	10	10	10
Moulds and Yeasts	≤100 CFU/g	100	100	100	100	100
<i>Salmonella</i>	Absent in 100 g	Negative	Negative	Negative	Negative	Negative
<i>Enterobacteriaceae</i>	Absent in 10 g	Negative	Negative	Negative	Negative	Negative
<i>Cronobacter spp.</i> (<i>Enterobacter sakazakii</i>)	Absent in 100 g	Negative	Negative	Negative	Negative	Negative
<i>Listeria monocytogenes</i>	Absent in 25 g	Negative	Negative	Negative	Negative	Negative
<i>Bacillus cereus</i>	≤50 CFU/g	10	10	10	10	10
Residual endotoxins	≤10 EU/mg	0.006	0.011	0.020	0.034	0.026
Aflatoxin M1	≤0.2 µg/kg	NT	0.02 (LOQ)	0.02 (LOQ)	NT	NT

CFU: colony forming units; EU: endotoxin units; NT: not tested; LOQ: limit of quantification.

Table 6. Compositional batch analysis of 6'-SL manufactured with an updated fermentation media without soy peptone: microbial contaminants.

		Batch H	Batch I	Batch J
Parameter	Specification			
Microbial parameters:				
Aerobic plate count	≤1,000 CFU/g	10	10	10
Moulds and Yeasts	≤100 CFU/g	10 (LOD)	10 (LOD)	10 (LOD)
<i>Salmonella</i>	Absent in 100 g	Negative	Negative	Negative
<i>Enterobacteriaceae</i>	Absent in 10 g	Negative	Negative	Negative
<i>Cronobacter spp.</i> (<i>Enterobacter sakazakii</i>)	Absent in 100 g	Negative	Negative	Negative
<i>Listeria monocytogenes</i>	Absent in 25 g	Negative	Negative	Negative
<i>Bacillus cereus</i>	≤50 CFU/g	10	10	10
Residual endotoxins	≤10 EU/mg	0.0001563 (LOQ)	0.0001563 (LOQ)	0.0001563 (LOQ)
Aflatoxin M1	≤0.2 µg/kg	0.02 (LOQ)	0.02 (LOQ)	0.02 (LOQ)

CFU: colony forming units; EU: endotoxin units; NT: not tested; LOQ: limit of quantification; LOD limit of detection.

21. The analysis parameters included purity, other carbohydrates, residual proteins, microbial parameters, heavy metals, sodium, and physicochemical properties (Annex B; p 15 – 18, Table 3.a-3 of response letter). A purity of 87% to 95% (dry weight basis (dwb)) is demonstrated by the analysis. The applicant reports a lack of heavy metals and residual proteins (≤ 100 mg/kg) and microbial parameters were within limits.

22. Other saccharide molecules N acetylneuraminic acid ($\leq 5.1\%$), D lactose ($\leq 0.05\%$), D glucose (0.05%), 6'-Sialyllactulose ($\leq 0.5\%$) were detected in low levels and 3'-Sialyllactose sodium salt was not detected. The applicant states that these molecules are natural components or break down products of human breast milk (Annex A; p 38 of dossier).

23. Absence of the genetically modified production organism and no detectable residual DNA in 3 batches of the final product is verified by PCR analysis (Annex A; p 31 – 32 of dossier). Results can be found in Annex C [Annex F].

All methods and validation are found in (Annex C [Annex G]). The applicant changed their analytical method for residual proteins from an internal method to the Bradford assay (Annex B; p 15 of response letter - confidential).

Stability

24. The applicant has provided a range of analytical data which characterises and describes the stability of 6'-SL. All 6'-SL batches analysed for stability were produced using a soy peptone media and due to little change in the composition of the novel food with soy peptone excluded from the fermentation media, the results from these stability studies were considered relevant. All study reports are found in (Annex C [Annex J]). The applicant reports the results from a 6-month accelerated storage conditions test ($40 \pm 2^\circ\text{C}$; $75 \pm 5\%$ Relative Humidity) on 5 batches as shown in (Annex A; p 32 – 34 and Table 2.c.3.1-1 of dossier). The parameters chosen for measurement and monitoring of stability includes physicochemical and biochemical (purity, carbohydrates and water content). The results show that parameters remained stable within specifications and the increase in the isomerisation product 6'sialyllactulose remains at low levels within limits (Annex B; p 27 - 28 of response letter and Annex A; p 32 of dossier).

25. The applicant further demonstrates the stability of parameters of the final novel food product under a 12-month normal conditions ($25 \pm 2^{\circ}\text{C}$; $60 \pm 5\%$ Relative Humidity) storage test on 1 batch. Microbiological stability is reported with regards to water activity values which remained stable under both accelerated and normal conditions (Annex A; p 34 - 35 of dossier). The applicant states that microbial growth and toxin formation is unlikely due to the low water content and low water activity of the final product. (Annex A; p 35 of dossier, and Annex B; p 30 of response letter).

26. Following request for further information on the data that supports the proposed 3-year shelf life, the applicant provided details of an Arrhenius equation used and further data (Annex B; p 27 - 30 of response letter). The applicant further reports results from a 24-month stability test on 1 batch under normal storage conditions (Annex B; p 29 and Table 3.d-1 of response letter) and results from a microbiological stability test at 24 months on 1 batch under normal storage conditions (Annex B; p 30 and Table 3.d-2 of response letter). Parameters remained stable and indicate that no microbial growth or toxin formation was detected.

Specifications

27. The specification parameters for the novel food ingredient were assessed using internationally recognised methods or are otherwise determined using internally developed and validated methods - see Table 7 below for the specifications (Annex A; p 38 - 39 , Table 2.d-1 of dossier and Annex C [Annexes G and H]).

Table 7. Specifications for 6'-SL.

Description: 6'-SL is a white to off-white powder that is produced by a microbial process. 6'-SL is isolated by spray-drying.

Source:

Genetically modified strain of *Escherichia coli* W

Parameter

Specification

Physicochemical

Appearance	Powder
Colour	White to off-white
pH	4.0 - 9.0
Composition	
Purity (6'-SL)	≥ 82 (% dwb)
NeuAc	≤ 9 w/w %
D-glucose	≤ 3 w/w %
D-lactose	≤ 3 w/w %
6'-sialyllactulose	≤ 5 w/w %
3'-sialyllactose sodium salt	≤ 1 w/w %
Water	≤ 10.5 w/w %
Sodium	≤ 5.0 w/w %
Residual protein (internal method)	≤ 10 mg/kg (0.001%)
Residual protein (Bradford assay)	≤ 100 mg/kg (1)

Heavy metals:

Arsenic ≤ 0.2 mg/kg

Cadmium ≤ 0.2 mg/kg

Lead ≤ 0.2 mg/kg

Mercury ≤ 0.2 mg/kg

Iron ≤ 10 mg/kg

Microbial parameters:

Aerobic plate count $\leq 1,000$ CFU/g

Moulds ≤ 100 CFU/g

Yeasts ≤ 100 CFU/g

Salmonella Absent in 100 g

Enterobacteriaceae Absent in 10 g

Cronobacter spp.
(*Enterobacter sakazakii*) Absent in 100 g

<i>Listeria monocytogenes</i>	Absent in 25 g
<i>Bacillus cereus</i>	≤50 CFU/g
Residual endotoxins	≤10 EU/mg
Aflatoxin M1	≤0.2 µg/kg

6'-SL: 6'-sialyllactose sodium salt; NeuAc: N acetylneuraminic acid; CFU: colony forming units; dwb: dry weight basis; EU: endotoxin units. (1) Evaluated using a test limit at 100 ppm.

History Of Use

28. The applicant conducts a literature search and review on the human biology of background exposure of 6'-SL from human breast milk and discusses likely intake levels based on the history of consumption of human breast milk in infants (Annex A; p 40 – 43 dossier). The applicant reports a GRAS notification in the USA for 6'-SL and one EU authorisation since 2021 (Annex A; p 43 of dossier).

29. The applicant conducts a review on the human biology of background exposure of 3'-SL and N-acetylneuraminic acid (NeuAc) from human breast milk and discusses intake levels based on the consumption of human breast milk (Annex A; p 43 – 45 of dossier). The applicant reports three GRAS notifications for 3SL in the USA, one GRAS notification for NeuAc, one EU authorisation for 3SL and further authorisations for NeuAc in the EU since 2021.

30. The applicant summarises the history regarding safety of the production microorganism (Annex A; p 40 of dossier). Details on the microorganism, applicable to a Category 1 food product within the GMO regulations are provided. As there is no presence of the microorganism in the final food it is not subject to review under the GMO regulations for food and feed Annex C [Annex B].

Proposed Use and Intake

31. The applicant states that the novel food ingredient is intended to be used by the general population including infants, children and individuals on energy

restricted diets for weight reduction.

32. Proposed food uses and maximum use levels are presented in Table 8 below (Annex A; p 46 – 47 , Table 2.f.2-1 of dossier). The applicant states that the proposed food uses and maximum use levels are the same as an existing 2020 EU novel food authorisation (Annex C – References – EFSA NDA Panel 2020a).

Table 8. Proposed uses and maximum use levels for 6'-SL.

EU Category Number	Category Name	Proposed Maximum use Level
1	Dairy products and analogues	
1.1	Unflavoured pasteurised and unflavoured sterilised (including UHT) milk	0.5 g/L
1.2/1.3	Unflavoured fermented milk-based products	0.5 g/L beverages 2.5 g/kg for products other than beverages
1.4	Flavoured fermented milk-based products including heat treated products	0.5 g/L beverages 5 g/kg for products other than beverages
7	Bakery wares	
7.2	Fine bakery wares (cereal bars only)	5 g/kg
13	Foods for Special Groups (FSG)	

13.1	Foods for infants and young children	
13.1.1	Infant formula as defined in Regulation (EU) No 609/2013	0.4 g/L in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer
13.1.2	Follow-on formula as defined in Regulation (EU) No 609/2013	0.3 g/L in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer
13.1.3	Processed cereal-based food and baby food for infants and young children as defined in Regulation (EU) No 609/2013	0.3 g/L in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer 2.5 g/kg for products other than beverages
13.1.4	Milk-based drinks and similar products intended for young children	0.3 g/L in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer
13.2	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	
13.2	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	On a case-by-case basis

13.3	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013	
		1.0 g/L beverages (equivalent to 0.25 g/meal based on a standard 250 g meal replacement beverage)
13.3	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013	10 g/kg for products other than beverages (equivalent to 0.30g/meal based on a standard 30 g meal replacement bar)
14	Beverages	
14.1.4	Flavoured drinks (excluding cola-type drinks)	0.5 g/L
17	Food supplements as defined in Directive 2002/46/EC	
		1.0 g/day for individuals above 3 years of age
17	Food supplements as defined in Directive 2002/46/EC	0.3 g/day for young children (12 to 35 months)
		0.4 g/day for infants (≤ 11 months)

6'-SL: 6'-sialyllactose sodium salt; UHT: ultra-high temperature.

33. The applicant refers to the estimations of mean and high anticipated intakes previously evaluated by EFSA's NDA Panel in 2020 (Annex A; p 47 - 48 of dossier). A summary is provided in Table 9 below (Annex A; p 48 , Table 2.f.3.1-1 of dossier).

Table 9. Mean and High Anticipated Intakes of 6'-SL from Proposed Food Uses Based on Summary Statistics from the Individual Food Intake Data from the EFSA Comprehensive Food Consumption Database (EFSA NDA Panel, 2020a)

Age groups	Number of EU Dietary Surveys	Range of Means (lowest and highest) among EU Dietary Surveys (mg/kg bw)	Range of 95th Percentile (lowest and highest) among EU Dietary surveys (mg/kg bw)
Infants (up to 11 months)	11	16 to 74	38 to 192
Young children or toddlers (12 to 35 months)	14	15 to 46	39 to 147
Other children (3 to 9 years)	19	6 to 25	16 to 61
Adolescents (10 to 17 years)	18	3 to 8	7 to 19
Adults (18 to 64 years)	19	1 to 5	5 to 13
Elderly (≥ 65 years)	18	1 to 4	5 to 10
Pregnant women	2	1 to 5	4 to 11
Lactating women	2	4 to 5	10 to 11

6'-SL: 6'-sialyllactose sodium salt; UHT: ultra-high temperature.

34. The applicant reports the highest estimated daily intake of 6'-SL amongst all population groups and all proposed uses, is in infants where at a maximum intake level 0.4 g/L is 104 mg/kg body weight/day, with the assumption that only infant formula would be consumed (Annex A; p 47 of dossier). The applicant reports the highest estimated daily intake, at the 95th percentile, based on proposed uses and maximum use levels in foods combined and taken from Food Intake Data from the EFSA Comprehensive Food Consumption Database, is 192 mg/kg body weight/day in infants (Annex C - References - EFSA NDA Panel 2020a). The applicant reports the estimated highest daily intake of 6'-SL from food intake alone is in breastfed infants at 193 mg/kg body weight/day.

35. The applicant states that proposed intakes are substitutional alternatives to other sources (breast milk or authorised products containing 6'-SL) rather than combined. Food supplements are not intended to be consumed with other sources on the same day. The applicant concludes from the assessment that the proposed intakes of 6'-SL are unlikely to exceed the estimated highest daily intake of 6'-SL in breastfed infants of 193 mg/kg body weight/day (Annex A; p 48 - 49 of dossier).

Absorption, Distribution, Metabolism, Excretion (ADME)

36. The applicant reports that the novel food ingredient 6'-SL is structurally and chemically identical to the naturally occurring counterpart in human milk and is considered a non-digestible oligosaccharide. The applicant refers to the EFSA NDA Panel 2020 evaluation and reports limited digestion in the upper gastrointestinal tract, limited absorption, fermentation by intestinal microbiota in the colon and 40 - 97% is excreted unchanged in faeces and 4% in urine. The ADME of the novel food ingredient is not expected to differ from the ADME of 6'SL from breast milk. (Annex A; p 50 - 51 of dossier).

37. The applicant states that the low levels of additional carbohydrate molecules in the final product consist of, those naturally present in breast milk, breakdown products of lactose and the isomerisation product 3'-SL whereby the overall exposure to these is expected to be insignificant. The applicant reports there are no laxative concerns regarding the low levels of 6'-sialyllactulose.

38. The applicant reports the results of a recent 2020 study on the ADME of 3'-SL and concludes that the ADME of 6'-SL and 3-SL are comparable and are constitutional isomers that possess minimal structural differences and have

similar functions and biological roles (Annex C – References – Galuska et al 2020).

Nutritional Information

39. The applicant refers to the EFSA NDA Panel 2020 evaluation that reports 6'-SL as not nutritionally disadvantageous.

40. 6'-SL is a non-digestible carbohydrate and fibre-type oligosaccharide (Annex A; p 52 of dossier).

41. The applicant refers to nutrient composition with regards to the low levels of additional carbohydrates in the final product composition and that overall exposure to these is expected to be insignificant (Annex A; p 52 of dossier). The applicant reports that the low levels of exposure to 6'-sialyllactulose in the final product (0.5%) is not expected to cause a laxative effect (Annex A; p 52 of dossier).

42. The novel food is expected to contribute to a low-level intake of sodium, with the final product containing $\leq 5\%$ sodium. The applicant reports on the highest estimated sodium intakes under proposed conditions of use for all population subgroups where the highest contribution is reported to be 24% of the safe daily intake in infants. The highest estimated intakes across all age groups from the proposed uses are presented in Table 2.h.2-1 (Annex A; p 53 - 54 , Table 2.h.2-1 of dossier).

43. The compositional batch analyses reported by the applicant detected D-lactose in low levels at $\leq 0.05\%$ (dry weight basis (dwb)) and a specification limit at $\leq 3\%$ is set by the applicant (Annex B; p 16, Table 3.a-3 of response letter).

Toxicological Information

44. The applicant conducted a bacterial reverse mutation assay (OECD 471) in accordance with GLP principles and reports that 6'-SL is non-mutagenic at concentrations of up to 5,000 $\mu\text{g}/\text{plate}$, in the absence or presence of metabolic activation (Annex A; p 57 of dossier). The full study report is available in Annex C [Annex L].

45. The applicant conducted an in vivo micronucleus test in mice (OECD 474) in accordance with GLP principles and reports that 6'-SL is non-genotoxic and does not induce chromosomal aberrations in oral doses of up to 2,000 mg 6'-SL/kg body weight (Annex A; p 58 of dossier). The full study report is available in Annex

C [Annex L].

46. The applicant conducted a 90-day oral toxicity study in rats (OECD 408) in accordance with GLP principles and reports a NOAEL of 2,168 mg/kg body weight/day for 6'-SL (Annex A; p 59 - 61 of dossier). The full study report is available in Annex C [Annex L] and a summary of statistically significant observations can be found in Annex C [Appendix B.3].

47. The applicant refers to the results of toxicological studies identified in the literature on comparable 6'SL preparations (Annex A; p 65, Table 2.i.3.1-1 of dossier) reporting comparable genotoxicity (Annex A; p 66 - 67 of dossier, Table 2.i.3.2-1) and subchronic toxicity findings (Annex A; p 68 - 69 of dossier, Table 2.i.3.3-1). The applicant refers to the results of two 21-day gastrointestinal developmental toxicity studies of 6'-SL in piglets that report no compound-related adverse effects in oral doses of 6'-SL of up to 1,200 mg 6'-SL/kg body weight/day (Annex A; p 71 - 72 of dossier, Table 2.i.3.5-1).

48. The applicant refers to results of toxicity testing conducted by the applicant on 3'-SL (Annex A; p 74- 79 of dossier). The applicant reports the results of a bacterial reverse mutation assay (OECD 471), an in vivo micronucleus test in mice (OECD 474), a 90-day oral toxicity study in rats (OECD 408) and two 21-day gastrointestinal developmental toxicity studies in piglets. The applicant reports that 3'-SL is non-mutagenic at concentrations of up to 5,000 µg/plate, in the absence or presence of metabolic activation, non-genotoxic and not inducing chromosomal aberrations in oral doses of up to 2,000 mg 3'-SL/kg body weight in mice, a NOAEL of >2,007 mg/kg body weight/day in rats and no compound-related adverse effects in piglets in oral doses of up to 1,200 mg 3'-SL/kg body weight/day, respectively. They suggest this indicates that there is not a concern from low levels of 3'-SL in the novel ingredient.

49. The applicant reports no safety concerns regarding any antimicrobial susceptibility or antimicrobial production with the proposed genetically modified culture microorganism (Annex A; p 105-107 of dossier). Results are reported from a phenotypic analysis (Annex C [Annex M]), genotypic analysis and genome search (Annex C [Annex C and Annex F]) and the applicant concludes absence of acquired antimicrobial resistance or potential for production of antimicrobial compounds.

Allergenicity

50. The applicant states that the high purification steps (microfiltration and ultra-filtration) in the production process demonstrate the lack of allergenic potential from any residual proteins ≤ 1 mg/kg (internal method) and ≤ 100 mg/kg (Bradford assay using a limit test at 100 ppm, Annex B; p 16 - 18 , Table 3.a.3 of response letter) and absence of the production strain in the final product (Annex A; p 108 , p 34 - 35 of dossier).

51. Lactose is added into the production process as a fermentation aid and is also a resulting minor component of the novel ingredient produced by fermentation itself. To exclude the potential for contamination of milk protein allergens, the applicant reports the results of two ELISA tests that did not detect milk proteins in the final 6'-SL product (Annex A; p 108 of dossier and Annex C [Annex F]). The applicant states that the altered nutrient media following the removal of soy peptone, removes the potential for soy allergens to be present (Annex A; p 16 - 20 and Table 2.b.1.2-1 of dossier) (Annex C [Annex D]) (Annex B; p 3 of response letter).

52. The applicant conducted a literature search and found no reports of sensitisation, case reports of allergic reactions, or allergenicity studies on 6'-SL (Annex A; p 109 of dossier and Annex C [Annex K]).

53. The applicant reports that the allergenic potential of introduced proteins as a result of the genetic modification to the culture microorganism were assessed using the National Institute of Health Sciences (Japan) Allergen Database for Food Safety (ADFS) (Annex A; p 108 of dossier). The applicant concludes that no sequence alerts for potential allergenicity were identified - Annex C [Annex N]. The culture microorganism is not present in the final ingredient including absence of any residual DNA.

Committee Action Required

- The Committee is asked whether the available data provide a satisfactory basis for evaluating the safety of this novel food ingredient.
- If so, the Committee is asked whether it is content to recommend approval for 6'-SL as a novel food ingredient to be added to the range of foods specified. A draft Committee Advice Document has been prepared and members comments are sought on the draft.
- If not, the Committee is asked to indicate what additional data would be required.

ACNFP Secretariat

September 2023

Annexes

Annex A - Dossier [Confidential]

Annex B - Request for information with applicant's responses [Confidential]

Annex C - Supporting documents with annexes, references and appendices
[Confidential]

Annex D - Draft Committee Advice Document